

248 Liver function and vitamin D in children with cystic fibrosis

M.F. Roddy^{1,2}, B. Elnazir², P. Grealley². ¹The National Childrens Hospital, Dietetics, Dublin, Ireland; ²AMNCH, Cystic Fibrosis Department, Dublin, Ireland

Introduction: The liver is important in the metabolism of vitamin D especially its hydroxylation to 25 OH D3. Liver disease occurs in up to 15–20% of children with CF raising the possibility that Vitamin D levels may be influenced by their liver function. Our aim was to look at vitamin D levels in children and to correlate it with their liver function tests.

Methods: A retrospective review of Vitamin D levels and liver function tests (AST, ALT, GGT, Alk Phos, Protein and Albumin) was conducted. Seventy-seven children with mean age of 12 were included in study. Minitab Statistical Package was used to analyse the data. Regression analysis and independent t tests were carried out to determine any relationship between vitamin D and liver function.

Results: Eleven children (14%) had a diagnosis of CF related liver disease (CFLD). The mean vitamin D was 73.12 (± 28 SD) with no difference in mean vitamin D between CFLD patients and none CFLD patients (mean of 65 nmol/ L versus 75 nmol/ L respectively, $p=0.27$). Vitamin D negatively correlated with GGT, protein and ALT ($p=0.02$, 0.04 and 0.02 respectively). Vitamin D did not correlate with albumin, AST or Alk Phos ($p=0.93$, 0.68 and 0.56 respectively). When those with CFLD were removed from the data, vitamin D still negatively correlated with protein ($p=0.04$) and GGT ($p=0.03$).

Conclusion: The cause of low vitamin D in CF is multifactorial. This study provides some evidence of a direct relationship (either cause or effect) between vitamin D and liver function. Further research is needed to determine if increased vitamin D supplementation in CF would affect liver function.

249 Influence of calcium and vitamin D on bone mineral density in children with cystic fibrosis

T. Jakovska-Maretti¹, S. Fustik¹, L. Spirevska¹. ¹University Pediatric Clinic, CF Center, Skopje, Macedonia, the Former Yugoslav Republic of

Low bone mineral density (BMD) is relatively frequent finding among patients with cystic fibrosis (CF). This complication is related with age, severity of lung damage and nutrition disorders.

Aim: To evaluate the effect of calcium and vitamin D supplementation on low BMD in patients with CF.

Materials and Methods: Z-score of BMD was assessed in 47 patients with CF and was performed with Quantitative Ultrasound. Chest deformation was found in 1/3 of schoolchildren with CF. Children with CF were divided into two groups: first group: 26 children younger than 12 years; second group: 21 children older than 12 years. All patients were pancreatic insufficient and received adequate basic therapy including multivitamins. Calcium and vitamin D supplements were given to correct the low values of BMD in children with CF for six months.

Results: Initial Z-score showed decreased values in both study groups: -0.14 ± 1.17 SD in CF children <12 years and -1.29 ± 0.9 SD in CF patients >12 years. 25OHD values in the younger group were 29.3 ± 15.2 ng/ml and 18.07 ± 8.3 ng/ml in older group. Calcium values were similar in the two groups (2.4 ± 0.1 ; 2.37 ± 0.2). After six months of treatment with calcium and vitamin D supplementation, the control densitometry showed significant increase in Z-score to 0.9 ± 1.5 SD in children <12 years and to -0.9 ± 1.1 SD in older CF patients.

Conclusion: Calcium and vitamin D have positive effect on mineral bone density in children with CF.

250 Vitamin D deficiency and thoracic kyphosis in adult cystic fibrosis patients

L. O'Shaughnessy¹, C. Reilly², E. Caples³, E. McKone⁴, C. Gallagher⁴. ¹St. Vincents University Hospital, Dietetics, Dublin, Ireland; ²St Vincent's University Hospital, Department of Physiotherapy, Dublin, Ireland; ³St Vincent's University Hospital, Department of Nutrition and Dietetics, Dublin, Ireland; ⁴St Vincent's University Hospital, National Referral Centre for Adult Cystic Fibrosis, Dublin, Ireland

Objectives: This study tested the hypothesis that there is an association between vitamin D deficiency and thoracic kyphosis in adults with Cystic Fibrosis (CF).

Methods: Subjects were recruited during annual review at the National Referral Centre for Adult Cystic Fibrosis. Each subject had thoracic curvature measured using a flexible ruler (previously validated against lateral x-ray), and had venous blood sampling to measure vitamin D level.

Results: 36 patients were included in the study, M:F ratio = 1.64:1.

5.5% of patients were in the severe lung disease category while 44% and 50% had moderate and mild disease respectively. 40.5% of patients were kyphotic (thoracic curvature greater than 40°). 13.8% of patients had optimal vitamin D levels (>75 nmol/L) and 2.7% were severely deficient with a measurement <25 nmol/L. A weak correlation of $r=0.17$ was found between angle of kyphosis and vitamin D levels. 43% of those patients studied had exposure to oral steroids.

Conclusions: The prevalence of vitamin D deficiency in this population was very high. But only weak correlation was found between the angle of thoracic kyphosis and vitamin D level. Further investigation would be of benefit to look at bone density and nutritional status in this population.

251 Should %ideal body weight (%IBW) or body mass index percentile (BMIp) be used to assess growth in children with cystic fibrosis?

K. Maslin¹, T. Curbishley¹, H. Kay², J. Legg². ¹University Hospital Southampton, Nutrition and Dietetics, Southampton, United Kingdom; ²University Hospital Southampton, Paediatric Respiratory Unit, Southampton, United Kingdom

Background: There is strong evidence that BMIp predicts nutritional failure more sensitively and accurately than %IBW, having a stronger correlation with %FEV1. This study aims to determine which growth measurement correlates most closely with %FEV1 in a UK population. Studies comparing both methods have concluded there is little difference for children of average height, but in short and tall children, underweight can be misclassified. %IBW is the main growth outcome measure in the 2002 CF Trust guidelines, which are currently under review.

Methods: Data was obtained from the 2010 South & South West regional CF database. Height, weight, BMIp and %FEV1 were calculated at annual review. BMIp was converted to Standard Deviation Scores (SDS) using UK reference data.

Results: Data was collected from 263 children (age range 5–17 years). Median %IBW was 101 (SD ± 16.17 , range 67–198). BMI SDS range was -3.21 to 3.57 (median -0.08 ± 1.13). Median %FEV1 was 87 (SD ± 19.8 , range 28–129). Overall, both %IBW and BMI SDS had a significant association with %FEV1, with BMI SDS having a stronger correlation ($r=0.3$, $p<0.01$). In children between 5–12 years old ($n=115$), the association between BMI SDS and %FEV1 was weaker ($r=0.18$, $p<0.05$), with no significant association between %IBW and %FEV1. In children with height >75 th percentile ($n=43$), neither growth measurement was significantly associated with %FEV1.

Conclusion: Compared to %IBW, BMIp is more closely associated with lung function in children <75 th percentile for height. The lack of correlation with either growth measurement and %FEV1 in tall children raises the possibility that other growth measures should be employed in this group.